

## EDITORIAL: REFLECTIONS ON PLANT CELL CLASSICS

### Genomic Balance Plays Out in Evolution

I will never forget the “Ah-ha” moment I had upon reviewing the manuscript by Blanc and Wolfe (2004) at the beginning of my stint as a co-editor of *The Plant Cell*. In the few years preceding that submission, it had become increasingly known from genome sequences that many species had evidence of whole genome duplications (WGD) in their evolutionary history. In *Arabidopsis*, there was evidence for at least three successive WGDs. Genes were present in duplicate in syntenic arrangements indicating that the two copies were not the result of local multiplication. Most genes, however, had returned to a singleton state to produce a near diploid gene repertoire. The contribution of Blanc and Wolfe was that the retained genes were not random in terms of their functions. As a general rule, transcription factors and signal transduction components were overrepresented among the duplicates present.

Our laboratory had a long standing interest in dosage sensitivity dating from the finding that changes in copy number of parts of genomes (aneuploidy) had a greater modulation of gene expression than changing the whole set of chromosomes (ploidy) (see Birchler and Veitia, 2007) suggesting that the stoichiometry of the genome had an impact on gene expression and quantitative characteristics (Birchler et al., 2001). Over the years we had sought single genes in *Drosophila* that might mimic the effects of aneuploidy giving inverse or positive correlative dosage modulations of gene expression across the genome. When the collection of these dosage sensitive modifiers was summarized, the functions of these genes were various transcription factors and signal transduction components (Birchler et al., 2001). Thus, there was a parallel between the classes of genes that were dosage sensitive on gene expression and the phenotype and those that were preferentially retained following WGD. Blanc and Wolfe noted this parallel.

Others in the field of evolutionary genomics came to a similar conclusion about the classes of genes that were retained following WGD (see Freeling, 2009). If the parallels to the aneuploidy/ploidy comparisons held, then there would be a complementary underrepresentation of the types of genes that were retained after WGD. In other words, if deletion of dosage sensitive genes were detrimental following WGD and selected against, then small scale duplications of the same classes of genes would also be detrimental in analogy to the behavior of plant aneuploids (Birchler and Veitia, 2007). This predication was in fact supported. A comprehensive analysis of retained genes following WGD and comparisons to representation in small scale duplications from sequence data across angiosperms reveals the generality of dosage balance sensitivity (Tasdighian et al., 2017).

The field has expanded to examine many aspects of WGD and the parameters of genome fractionation. An interesting phenomenon discovered from these studies is the phenomenon of subgenome dominance (Freeling, 2009). When divergent genomes come together to form an allopolyploid, the two subgenomes that result often have different levels of overall expression. This difference is maintained over evolutionary time and can have an impact on genome fractionation. In a study of subgenome dominance in monkeyflower, a comparison was made of an interspecific hybrid, a newly produced allopolyploid and a natural allopolyploid to compare this difference (Edger et al., 2017). An interplay of transposon density in subgenomes, DNA methylation and gene expression is implicated in this dominance.

Micro RNAs are small RNAs that often modulate the levels of transcription factor expression by their impact on target mRNA levels. When the miRNA genes were examined in the grass family, it was found that duplicates were over-retained following WGD and that there was a differential fractionation between the subgenomes (Abrouk et al., 2012). There is a co-retention of miRNAs and their targets indicating a resistance to differential return to the singleton state and a need to maintain the regulatory balance.

Further elaboration of fractionation involved an examination of the metabolic network of *Arabidopsis* (Bekaert et al., 2011). These authors studied the interplay between relative dosage of duplicates and absolute dosage. They note that gene duplication will be maintained for enzymes for which a high copy number is needed to maintain metabolic flux. They identified a difference in flux bias between the two most recent WGDs and discussed how relative versus absolute dosage in the resolution of duplicates occurs following WGD.

The retention of duplicate genes involved in regulatory processes for longer periods of evolutionary time provides the grist for evolutionary innovation (Birchler and Veitia, 2012). While such a contention might appear contradictory given the selection pressure to maintain a balanced dosage, it is clear that new functional balances can emerge (Birchler and Veitia, 2007) as evidenced, for example, by the fact that of the three readily discernible WGDs in the *Arabidopsis* genome, there is a decreasing number of retained genes with receding time from the present. As new functional balances evolve, new morphological innovations can emerge and the duplicate pairs can undergo sub or neo functionalization that will lock their duplicate nature in place going forward.

The role of genomic balance in evolution is an ongoing active area of research. There are many aspects of the role of “balance” in expression (Hou et al., 2018) and evolution that are exciting avenues for future exploration. Understanding the interplay of networks and regulatory cascades and the

emergence of new functional balances over time has been little explored to date but will become more accessible with new sequencing technologies. Coupling these data with biochemical studies of assembly and interactions of multisubunit regulatory machines is needed to determine the critical steps molecularly at the cellular level that condition the retention of dosage sensitive duplicates in populations. Intercalating data on global gene expression with population and evolutionary genomics of regulatory genes will help elucidate what stoichiometric relationships impact reproductive fitness to varying degrees. The pages of *The Plant Cell* have included many seminal papers in the establishment of this field and hopefully will continue to do so in the future.

**James A. Birchler**  
Division of Biological Sciences  
University of Missouri  
Columbia, MO 65211  
Email: [BirchlerJ@Missouri.edu](mailto:BirchlerJ@Missouri.edu)  
ORCID: 0000-0003-3643-2756

## REFERENCES

- Abrouk, M., Zhang, R., Murat, F., Li, A., Point, C., Mao, L., and Salse, J.** (2012). Grass microRNA gene paleohistory unveils new insights into gene dosage balance in subgenome partitioning after whole genome duplication. *Plant Cell* **24**: 1776-1792.
- Bekaert, M., Edger, P.P., Pires, J.C., and Conant, G.C.** (2011). Two-phase resolution of polyploidy in the Arabidopsis metabolic network gives rise to relative and absolute dosage constraints. *Plant Cell* **23**: 1719-1728.
- Birchler, J.A., Bhadra, U., Pal Bhadra, M., and Auger, D.L.** (2001). Dosage dependent gene regulation in multicellular eukaryotes: Implications for dosage compensation, aneuploid syndromes and quantitative traits. *Developmental Biology* **234**: 275-288.
- Birchler, J.A. and Veitia, R.A.** (2007). The gene balance hypothesis: from classical genetics to modern genomics. *Plant Cell* **19**: 395-402.
- Birchler, J.A. and Veitia, R.A.** (2012). Gene balance hypothesis: connecting issues of dosage sensitivity across biological disciplines. *Proc. Natl. Acad. Sci. USA* **109**: 14746-14753.
- Blanc, G. and Wolfe, K.H.** (2004). Functional divergence of duplicated genes formed by polyploidy during Arabidopsis evolution. *Plant Cell* **16**: 1679-1691.
- Edger, P.P., Smith, R., McKain, M.R., Cooley, A.M., Vallejo-Marin, M., Yuan, Y., Bewick, A.J., Ji, L., Platts, A.E., Bowman, M.J., Childs, K.L., Washburn, J.D., Schmitz, R.J., Smith, G.D., Pires, J.C., and Puzey, J.R.** (2017). Subgenome dominance in an interspecific hybrid, synthetic allopolyploid, and a 140-year-old naturally established neo-allopolyploid monkeyflower. *Plant Cell* **29**: 2150-2167.
- Freeling, M.** (2009). Bias in plant gene content following different sorts of duplication: tandem, whole-genome, segmental, or by transposition. *Annu. Rev. Plant Biol.* **60**: 433-453.
- Hou, J., Shi, X., Chen, C., Islam, M.S., Johnson, A.F., Kanno, T., Huettel, B., Yen, M.R., Hsu, F.-M., Ji, T., Chen, P.-Y., Matzke, M., Matzke, A.J.M., Cheng, J., and Birchler, J.A.** (2018). Global impacts of chromosomal imbalance on gene expression in Arabidopsis and other taxa. *Proc. Natl. Acad. Sci. USA* **115**: 11321-11330.
- Tasdighian, S., Van Bel, M., Li, Z., Van de Peer, Y., Carretero-Paulet, L., and Maere, S.** (2017). Reciprocally retained genes in the angiosperm lineage show the hallmarks of dosage balance sensitivity. *Plant Cell* **29**: 2766-2785.

## Genomic Balance Plays Out in Evolution

James A. Birchler

*Plant Cell*; originally published online May 2, 2019;  
DOI 10.1105/tpc.19.00329

This information is current as of June 10, 2019

|                                 |   |
|---------------------------------|---|
| <b>Permissions</b>              | <a href="https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;issn=1532298X&amp;WT.mc_id=pd_hw1532298X">https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;issn=1532298X&amp;WT.mc_id=pd_hw1532298X</a> |
| <b>eTOCs</b>                    | Sign up for eTOCs at:<br><a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>  |
| <b>CiteTrack Alerts</b>         | Sign up for CiteTrack Alerts at:<br><a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>   |
| <b>Subscription Information</b> | Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at:<br><a href="http://www.aspb.org/publications/subscriptions.cfm">http://www.aspb.org/publications/subscriptions.cfm</a>        |